arXiv:math/0502426v1 [math.RA] 20 Feb 2005

The group of automorphisms of the category of free associative algebras

A. Berzins
University of Latvia
e-mail: aberzins@latnet.lv

August 2004

AMS Mathematics Subject Classification: 14A99, 14P05
Keywords: Associative algebra, automorphism, variety

Abstract

In this paper, the problem formulated in [8] is solved. We prove, that the group of automorphisms of the category of free associative algebras is generated by semi-inner and mirror automorphisms.

1 Introduction

The classical algebraic geometry arise from consideration systems of polynomial equations in an affine space P^n, where P is arbitrary field. Although equations solving in algebras in different varieties of algebras Θ was considered long ago, foundation of the universal algebraic geometry are B. Plotkin's scientific works [5-8] where concepts of affine space, algebraic set, algebraic variety for arbitrary varieties of algebras were introduced.

The basis of this theory is consideration of a point as an element from $\mathrm{Hom}(W, X)$, where W is a free finitely generated algebra, and X is a fixed algebra in variety of universal algebras Θ. It was found that many problems in this theory, such as geometric equivalence, geometric similarity, isomorphism and equivalence of categories of algebraic sets and varieties depend on structure of $\mathrm{Aut}(\Theta^0)$ and $\mathrm{Aut}(\mathrm{End}(W))$, where Θ^0 is the category of algebras

$W = W(X)$ with the finite X that are free in Θ, and W is a free algebra in Θ. Structure of $\mathrm{Aut}(\mathrm{Com}-P))^0$ (i.e, the classical case) was described in [1]. Also are described $\mathrm{Aut}(\Theta^0)$ for varieties of groups, semigroups, Lie algebras and some another varieties. For some varieties are described $\mathrm{Aut}(\mathrm{End}(W))$, but these problems are more difficult. In particular, this problem is not solved even in the classical case of free commutative algebras, i.e. the structure of $\mathrm{Aut}(\mathrm{End}(P[x_1,\dots,x_n]))$ is not describe if $n > 2$. In all solved cases groups $\mathrm{Aut}(\mathrm{End}(W))$ are generated by semi-inner automorphisms and mirror automorphism.

In this article we describe $\mathrm{Aut}(\mathrm{Ass}-P)^0$. When the paper had been finished, I was informed that a solution of this problem is contained in the forthcoming paper [4]. Both solutions of this problem are principally different, and methods of proves are independent. Explicitly I want to note that proof in the present paper does not use the description of the group $\mathrm{Aut}(W(x,y))$. It is very important, because structure of the group $\mathrm{Aut}(W(x_1,\dots,x_n))$ is not described, when $n > 2$, but counterexamples show that structures of $\mathrm{Aut}(W(x_1,\dots,x_n))$ and $\mathrm{Aut}(W(x,y))$ are principally different. So, methods presented in the paper may be useful to describe the group $\mathrm{Aut}(\mathrm{End}(W_n))$ for $n > 2$.

2 Definitions

Let $(\mathrm{Ass}-P)^0 = \Theta^0$ be the category of free associative algebras over a field P with a finite set of generators, and $\tau \in \mathrm{Aut}(\Theta^0)$. Then there exist a set of bijections $\mu = \{\mu_i\}$, $i \in \mathrm{N}$ of W_i, such that for every $s \in \mathrm{Hom}(W_i, W_j)$

$$s^\tau = \mu_i s \mu_j^{-1}$$

Here W_k is the free k-generated associative algebra. Automorphism τ which is representable in the form $s^\tau = \mu s \mu^{-1}$ is called quasi-inner. In the category $(\mathrm{Ass}-P)^0$ every automorphism is quasi-inner. Of course, arbitrary set of substitutions does not generate an automorphism. For category Θ^0 we have three types of bijections μ which generate an automorphism.

1. $\mu = \overline{\alpha}$, where $\overline{\alpha}$ is the natural extension of an automorphism $\alpha \in \mathrm{Aut}(P)$ on W, i.e. $\overline{\alpha}\left(\sum a_k u_k\right) = \sum \alpha(a_k) u_k$. We shall write $\overline{\alpha} = \alpha$ and call it "automorphism of field".Here u_k is a product of variables.

2. $\mu = \eta = \{\eta_i\}$, $\eta_i \in \mathrm{Aut}(W_i)$.

Let as recall some definitions for Ass-P. (See [8] for general case)

Definition 2.1 *The automorphism τ generated by a set of automorphisms $\eta = \{\eta_i \in W_i\}$ is called inner automorphism.*

Definition 2.2 *Note, that every $\alpha \in Aut(P)$ belongs to normalizer of subgroup $Aut(W_i)$ in the group of all bijections of W_i. So, every product of elements of $Aut(P)$ and $Aut(W_i)$ my be represented in the form $\mu_i = \alpha\eta_i$, $\alpha \in Aut(P)$, $\eta_i \in Aut(W_i)$. Automorphism $\tau \in Aut(\Theta^0)$ generated by μ is called semi-inner.*

3. Now we describe the third type of μ. Let $S = S(X)$ be a free semigroup. For every $u = x_{i_1}x_{i_2}\dots x_{i_n}$ in S, take $\overline{u} = x_{i_n}\dots x_{i_2}x_{i_1}$. Then $u \to \overline{u}$ is an antiautomorphism of the semigroup S.

Definition 2.3 *Let $W = W(X)$ be a free associative algebra. For every its element $\omega = \lambda_0 + \lambda_1 u_1 + \cdots + \lambda_k u_k$ denote $\overline{\omega} = \lambda_0 + \lambda_1\overline{u_1} + \cdots + \lambda_k\overline{u_k}$. The transition $\omega \to \overline{\omega}$ is an antiautomorphism of the algebra W. The set of these transitions $\beta = \{\beta_i : W_i \to W_i\}$ generate the automorphism $\delta \in Aut(\Theta^0)$, $s^\delta = \beta s \beta^{-1}$. We call β mirror antiautomorphism of W and δ mirror automorphism of $Aut(\Theta^0)$.*

Here δ is not inner and is not semi-inner, but is quasi-inner. Note that

1. δ belongs to the normalizer of subgroup $\mathrm{Sinn}(\Theta^0)$ in $\mathrm{Aut}(\Theta^0)$ that consists of all semi-inner automorphisms;
2. every antiautomorphism of W is a product of the mirror antiautomorphism β and an automorphism of W.

3 Almost central algebras

Let $W(X)$ be a free finitely generated algebra in variety Θ.

Definition 3.1 *Bijection $\mu : W(X) \mapsto W(X)$ is called central, if for every $s \in End(W(X))$*

$$\mu s = s\mu.$$

Algebra $W(X)$ is called central, if every its central bijection is identical.

Theorem 1 *Let $W(X)$ be a free central algebra and μ some bijection of $W(X)$ generating an automorphism $\tau \in Aut(End(W(X)))$. Then μ transforms every base of algebra $W(X)$ to a base of algebra.*

Proof. [see 3]

For free algebras in a variety Θ of algebras over field P (Com-P, Ass-P, and others) we shall correct the definition of central algebra. Clearly, in every such algebra $W(X)$ we have a bijection $\mu(u) = au + b$, where $a, b \in P$, $a \neq 0$ (linear bijection). This bijection commutate with every endomorphism of algebra $W(X)$ and, obviously, transforms every base of $W(X)$ to a base.

Definition 3.2 *Finitely generated algebra $W(X)$ over field P is called almost central, if every its central bijection is linear.*

Theorem 2 *Free finitely generated commutative and associative algebras are almost central.*

Proof. Let μ be a central bijection of $W(X)$, $\mu(x_i) = r_i(x_1, x_2, \ldots, x_n) \in W(X)$, and $s \in \mathrm{End}(W(X)$, $s(x_1) = x_1$, $s(x_i) = 0$ for $i = 1$. Then

$$\mu s(x_1) = \mu(x_1) = r_1(x_1, x_2, \ldots, x_n),$$

$$s\mu(x_1) = s(r_1(x_1, x_2, \ldots, x_n)) = r_1(x_1, 0, \ldots, 0) = r(x_1).$$

So $r_1(x_1, x_2, \ldots, x_n) = r(x_1)$ is a polynomial of one variable.

For arbitrary $u \in W(X)$ take $s \in \mathrm{End}(W(X))$ $s(x_1) = u$, $s(x_i) = x_i$ for $i > 1$. Then

$$\mu(u) = \mu s(x_1) = s\mu(x_1) = s(r(x_1)) = r(u),$$

and so $\mu(u) = r(u) = a_0 + a_1 + \cdots + a_k u^k$. Because μ is bijection, so r is linear, Q.E.D..

Corollary. From the theorems 1 and 2 we have that in mentioned algebras every bijection μ, which generate an automorphism of $\mathrm{Aut}(\mathrm{End}(W(X)))$ transforms every base of algebra $W(X)$ to a base of algebra.

4 The group $\mathrm{Aut}(\mathrm{Ass} - P)^0$

Now we can formulate the main result of this article (problem 6 in [8]).

Theorem 3 *The group* $Aut(Ass{-}P)^0$ *is generated by semi-inner and mirror automorphisms.*

Proof. Let τ be an automorphism of Θ^0. Then exist a set of bijections $\mu = \mu_i$, such that for every $s \in \mathrm{Hom}(W_i, W_j)$ equality $s^\tau = \mu_j s \mu_i^{-1}$ holds. We must prove that $\mu_i = \alpha \cdot \eta_i$ for every $i \in \mathrm{N}$ or $\mu_i = \alpha \cdot \eta_i \cdot \beta$ for every $i \in \mathrm{N}$. Here $\alpha \in \mathrm{Aut}(P)$, $\eta_i \in \mathrm{Aut}(W_i)$ and β is mirror.

Let us consider the free 1-generated associative algebra $W(x) = P[x]$, i.e. the free 1-generated commutative algebra. From the analogous theorem for $\mathrm{Aut}(\mathrm{Com} - P)^0$ (see [1]) we have that $\mu_1 = \alpha\eta_1$. So, we can consider $\mu' = \alpha^{-1}\mu$. Then μ'_1 is automorphism of $W(x)$. Clearly, the theorem will be proved, if we check that μ' is a set of automorphisms or a set of antiautomorphisms of W_i.

Write $\mu = \mu'$. Let $s \in \mathrm{Hom}(W_1, W_2)$, $s : t \to f(x, y)$. It follows from $\mu_2 s = s^\tau \mu_1$ that μ_2 is a morphism of 1-generated subalgebra $P(f(x, x))$ of $W(x, y)$ to $W(x, y)$.

Let $s \in S = \mathrm{Hom}(W(x, y), W(t))$. Then $\mu_1 s = s^\tau \mu_2$; $\mu_1 s(x) = s^\tau \mu_2(x)$, $\mu_1 s(y) = s^\tau \mu_2(y)$, $\mu_1 s(x + y) = s^\tau \mu_2(x + y)$. Denote $\mu = \mu_2$. Then

$$s^\tau(\mu(x + y) - \mu(x) - \mu(y) = \mu_1 s((x + y) - x - y) = 0,$$

and

$$\mu(x + y) - \mu(x) - \mu(y) \in \bigcap_{s\in S} \mathrm{Ker} s = (xy - yx) \tag{1}$$

Here $(xy - yx)$ is ideal generated by $xy - yx$. Analogously

$$\mu(x \cdot y) - \mu(x) \cdot \mu(y) \in (xy - yx) \tag{2}$$

Lemma 4.1 $\mu(u + v) = \mu(u) + \mu(v)$ *for every* $u, v \in W(X)$.

Proof. Let

$$\Delta(x, y) = \mu(x + y) - \mu(x) - \mu(y) = f(x, y) = g(\mu(x), \mu(y).$$

Last equality holds because by theorem 2 algebra $W(X)$ is almost central. Note that $g(u,v)$ is homogenous of degree 1. Really,

$$g(\alpha\mu(x), \alpha\mu(y)) = g(\mu(\alpha x), \mu(\alpha y)) =$$

$$\mu(\alpha x + \alpha y) - \mu(\alpha x) - \mu(\alpha y) = \alpha \cdot g(\mu(x), \mu(y)).$$

So $\mu(x+y)-\mu(x)-\mu(y) = a\mu(x)+b\mu(y)$, i.e. $\mu(x+y) = c\mu(x)+d\mu(y)$. Using endomorphism we have $\mu(u+v) = c\mu(u) + d\mu(v)$ for every $u, v \in W(X)$. Substitutions $u = 0$ and $v = 0$ gives that $c = d = 1$, Q.E.D.

Lemma 4.2 *$\mu(xy) = \mu(x)\mu(y) + a(xy - yx)$, where $a \in P$.*

Proof. Let $\Delta(x,y) = \mu(xy)-\mu(x)\cdot\mu(y)$. Then $\Delta(x,y)$ is x and y homogenous of degree 1. Since $\Delta(x,y) \in (xy-yx)$ (formula (2)), so $\Delta(x,y) = a(xy-yx)$. Q.E.D.

Lemma 4.3 *$\mu(xy) = \mu(y)\mu(x) + b(xy - yx)$, where $b \in P$.*

The proof is the same as the proof of the lemma 4.2.

Lemma 4.4 *$\mu(xy) = \alpha\mu(x)\mu(y) + \beta\mu(y)\mu(x)$, where $\alpha, \beta \in P$.*

Proof. It follows from lemma 4.2 and lemma 4.3 that

$$(b-a)\mu(xy) = b\mu(x)\mu(y) - a\mu(y)\mu(x)$$

If $a = b$, then from lemma 4.2 and lemma 4.3 we get equality $\mu(x)\mu(y) = \mu(y)\mu(x)$ which easy leads to contradiction. So we can take $\alpha = \frac{b}{b-a}$ and $\beta = \frac{-a}{b-a}$ and get required equality.

Corollary. For every $u, v \in W(x,y)$

$$\mu(uv) = \alpha\mu(u)\mu(v) + \beta\mu(v)\mu(u) \tag{3}$$

Lemma 4.5 *$\mu(uv) = \mu(u)\mu(v)$ for every $u, v \in W(x,y)$ or*
$\mu(uv) = \mu(v)\mu(u)$ for every $u, v \in W(x,y)$.

Proof. Let u, v be elements of $W(x,y)$, such that $\mu(u) = x$ and $\mu(v) = y$. Let us denote $\mu(r) = \overline{r}$. Then

$$\mu(uuv) = \alpha\overline{uu} \cdot \overline{v} + \beta\overline{v} \cdot \overline{uu} = \alpha xxy + \beta yxx,$$

$$\mu(uuv) = \alpha\overline{u} \cdot \overline{uv} + \beta\overline{uv} \cdot \overline{u} = \alpha^2 xxy + 2\alpha\beta xyx + \beta^2 yxx.$$

Therefore $\alpha^2 = \alpha$, $\alpha\beta = 0$, $\beta^2 = \beta$. Because the case $\alpha = \beta = 0$ is impossible, we have two solutions:

1. $\beta = 0, \alpha = 1$, then $\mu(uv) = \mu(u)\mu(v)$;
2. $\beta = 1, \alpha = 0$, then $\mu(uv) = \mu(v)\mu(u)$.

It follows from $\mu(\alpha u) = \alpha\mu(u)$, lemma 4.1 and lemma 4.5 that μ is an automorphism or antiautomorphism of $W(x,y)$. Using endomorphisms $s \in \mathrm{Hom}(W_2, W_n)$ it is easy to check that μ_n also is an automorphism or antiautomorphism. The theorem is proved.

Acknowledgments

The author is happy to thank professor B. Plotkin for stimulating discussions of the results.

References

1. A. Berzins, *The automorphisms of* End$K[x]$, Proc. Latvian Acad. Sci., Section B, 2003, vol. 57, no. 3/4, pp. 78-81.
2. A. Berzins, *Geometric equivalence of algebras*, Int. J. Alg. Comput., 2001, vol. 11, no. 4, pp. 447-456.
3. A. Berzins, B. Plotkin and E. Plotkin, *Algebraic geometry in Varieties with the Given Algebra of constants*, J. Math. Sci., New York, 2000, vol. 102, no. 3, pp. 4039-4070.
4. G. Mashevitzky, *Automorphisms of categories of free associative algebras*, Preprint.
5. B. Plotkin, *Algebraic logic, varieties of algebras and algebraic varieties*, Proc. Int. Alg. Conf., St. Petersburg, 1995, Walter de Gruyter, New York, London, 1996.

6. B. Plotkin, *Varieties of algebras and algebraic varieties*, Israel J. of Mathematics, 1996, vol. 96, no. 2, pp. 511-522.
7. B. Plotkin, *Varieties of algebras and algebraic varieties. Categories of algebraic varieties*, Sib. Adv. Math., 1997, vol. 7, no. 2, pp. 64-97.
8. B. Plotkin, *Algebras with the same (algebraic) geometry*, Proc. Steklov Inst. Math, Vol. 242, 2003, pp. 165-196.